

**Table 7.6f**  
**Immunosuppression Use for Maintenance by Regimen**  
**One Year Following Transplantation, 1997 to 2006**  
**Recipients with Pancreas After Kidney (PAK) Transplants**

	Transplant Year									
	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
<b>Transplants</b>	130	156	220	304	305	375	344	420	343	293
<b>Functioning Graft at 1 Year PostTx</b>	89	110	167	219	245	272	259	321	256	221
<b>With Maintenance Use Recorded at 1 Year PostTx</b>										
Tac	19	70	163	211	242	267	251	318	254	218
--- + MMF/MPA	0.0%	0.0%	0.0%	0.9%	0.0%	0.7%	1.6%	5.3%	3.1%	1.8%
CyA + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.4%	0.7%	0.8%	0.9%	1.2%	1.4%
Tac + MMF/MPA	0.0%	0.0%	1.2%	2.8%	4.1%	3.4%	7.6%	8.5%	18.5%	19.3%
Tac + OtherAntimet	0.0%	1.4%	0.0%	0.9%	0.0%	0.0%	0.8%	0.0%	0.4%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.6%	0.4%	2.3%
Tac + Siro/Evero	0.0%	0.0%	0.0%	0.9%	4.1%	7.5%	4.4%	7.2%	4.7%	4.6%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.8%	0.3%	1.6%	1.4%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.4%	1.2%	1.3%	2.4%	1.8%
CyA + OtherAntimet + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.4%	0.0%	0.0%	0.0%	0.0%	0.0%
--- + Steroids	0.0%	0.0%	0.0%	0.5%	0.0%	0.0%	0.4%	0.0%	0.0%	0.0%
CyA + Steroids	0.0%	1.4%	0.0%	0.9%	0.4%	0.4%	1.2%	0.0%	0.4%	0.0%
Tac + Steroids	5.3%	1.4%	3.7%	4.7%	4.1%	4.1%	3.2%	2.2%	0.8%	3.7%
--- + MMF/MPA + Steroids	0.0%	0.0%	0.6%	0.5%	0.4%	0.4%	0.0%	0.3%	0.4%	1.8%
CyA + MMF/MPA + Steroids	10.5%	22.9%	9.2%	8.1%	8.3%	9.7%	5.2%	4.1%	2.0%	2.8%
Tac + MMF/MPA + Steroids	73.7%	62.9%	73.0%	54.5%	55.4%	53.2%	45.8%	50.3%	54.3%	50.9%
--- + OtherAntimet + Steroids	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%
CyA + OtherAntimet + Steroids	0.0%	1.4%	0.6%	0.5%	0.0%	0.4%	0.0%	0.0%	0.4%	0.0%
Tac + OtherAntimet + Steroids	10.5%	4.3%	4.3%	2.4%	0.8%	0.4%	1.6%	0.9%	0.8%	1.8%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.6%	0.5%	0.8%	0.4%	0.4%	0.0%	0.0%	0.5%
CyA + Siro/Evero + Steroids	0.0%	1.4%	1.8%	2.8%	1.2%	1.9%	0.8%	0.6%	0.8%	0.5%
Tac + Siro/Evero + Steroids	0.0%	0.0%	3.7%	14.7%	15.3%	10.9%	10.0%	7.2%	4.3%	3.7%
Other Regimen	0.0%	2.9%	1.2%	4.3%	4.1%	5.2%	12.7%	6.6%	3.5%	1.4%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages are calculated based on the total number of patients with maintenance use at one year following transplantation. They sum to 100 percent for each column.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.